

# Mental Health Mediates the Association Between Gender-Based Violence and HIV Treatment Engagement in US Women

Amy A. Conroy, PhD, MPH,<sup>a</sup> Jennifer P. Jain, PhD,<sup>b</sup> Lila Sheira, MPH,<sup>c</sup> Edward A. Frongillo, PhD,<sup>d</sup> Torsten B. Neilands, PhD,<sup>a</sup> Mardge H. Cohen, MD,<sup>e</sup> Tracey E. Wilson, PhD,<sup>f</sup> Aruna Chandran, MD, MPH,<sup>g</sup> Adaora A. Adimora, MD, MPH,<sup>h</sup> Seble Kassaye, MD,<sup>i</sup> Anandi N. Sheth, MD, MSc,<sup>j</sup> Margaret A. Fischl, MD,<sup>k</sup> Adebola Adedimeji, PhD, MPH, MBA,<sup>l</sup> Janet M. Turan, PhD, MPH,<sup>m</sup> Phyllis C. Tien, MD,<sup>n</sup> and Sheri D. Weiser, MD, MPH<sup>a,c</sup>

**Background:** Gender-based violence (GBV) is associated with poorer engagement in HIV care and treatment. However, there is a dearth of research on the psychological (eg, mental health) and structural (eg, food insecurity) factors that mediate and moderate this association. GBV could lead to poor mental health, which in turn affects adherence, whereas food insecurity could worsen the effect of

GBV on engagement in care. This study uses data from the Women's Interagency HIV Study to address these gaps.

**Methods:** Women completed 6 assessments from 2013 to 2016 on GBV, mental health, food insecurity, adherence to antiretroviral therapy, and missed HIV care appointments in the past 6 months. Multilevel logistic regression models estimated associations between

Received for publication August 16, 2021; accepted October 11, 2021.

From the <sup>a</sup>Department of Medicine, Center for AIDS Prevention Studies, University of California, San Francisco, CA; <sup>b</sup>Department of Psychiatry and Behavioral Sciences, University of California, San Francisco, CA; <sup>c</sup>Division of HIV, Infectious Diseases, and Global Medicine, University of California, San Francisco, CA; <sup>d</sup>Department of Health Promotion, Education, and Behavior, University of South Carolina, Columbia, SC; <sup>e</sup>Department of Medicine, Stroger Hospital, Chicago, IL; <sup>f</sup>Department of Community Health Sciences, State University of New York Downstate Health Sciences University, School of Public Health, Brooklyn, NY; <sup>g</sup>Department of Epidemiology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, MD; <sup>h</sup>School of Medicine and UNC Gillings School of Global Public Health, University of North Carolina at Chapel Hill, Chapel Hill, NC; <sup>i</sup>Department of Family Medicine, Georgetown University Medical Center, Washington, DC; <sup>j</sup>School of Medicine, Emory University, Grady Health System, Atlanta, GA; <sup>k</sup>Department of Medicine, University of Miami, Miami, FL; <sup>l</sup>Department of Epidemiology and Population Health, Albert Einstein College of Medicine, Bronx, NY; <sup>m</sup>Department of Health Care Organization and Policy, School of Public Health, University of Alabama at Birmingham, Birmingham, AL; and <sup>n</sup>Department of Medicine, Department of Veteran Affairs Medical Center, UCSF and Medical Service, San Francisco, CA.

This study was funded by a Women's Interagency HIV Study (WIHS) substudy Grant from the National Institute of Mental Health, R01MH095683, and a mentoring Grant K24AI134326 (S.D.W.). Additional salary support for A.A.C. was provided by P30AI027763. J.P.J. is supported by a postdoctoral fellowship award program from the National Institute on Drug Abuse (T32DA007250). Data in this manuscript were collected by the Women's Interagency HIV Study (WIHS), now the MACS/WIHS Combined Cohort Study (MWCCS). The contents of this publication are solely the responsibility of the authors and do not represent the official views of the National Institutes of Health (NIH). MWCCS (Principal Investigators): Atlanta CRS (Ighovwerha Ofotokun, A.N.S., and Gina Wingood), U01-HL146241; Baltimore CRS (Todd Brown and Joseph Margolick), U01-HL146201; Bronx CRS (Kathryn Anastos and Anjali Sharma), U01-HL146204; Brooklyn CRS (Deborah Gustafson and T.E.W.), U01-HL146202; Data Analysis and Coordination Center (Gypsyamber D'Souza, Stephen Gange and Elizabeth Golub), U01-HL146193; Chicago-Cook County CRS (M.H.C. and Audrey French), U01-HL146245; Chicago-Northwestern CRS (Steven Wolinsky), U01-HL146240; Northern California CRS (Bradley Aouizerat, Jennifer Price, and P.C.T.), U01-HL146242; Los Angeles CRS (Roger Detels and Matthew Mimiaga), U01-HL146333; Metropolitan Washington CRS (S. Kassaye and Daniel Merenstein), U01-HL146205; Miami CRS (Maria Alcaide, M.A.F., and Deborah Jones), U01-HL146203; Pittsburgh CRS (Jeremy Martinson and Charles Rinaldo), U01-HL146208; UAB-MS CRS (Mirjam-Colette Kempf, Jodie Dionne-Odom, and Deborah Konkle-Parker), U01-HL146192; and UNC CRS (A.A.A.), U01-HL146194. The MWCCS is funded primarily by the National Heart, Lung, and Blood Institute (NHLBI), with additional cofunding from the Eunice Kennedy Shriver National Institute of Child Health and Human Development (NICHD), National Institute on Aging (NIA), National Institute of Dental and Craniofacial Research (NIDCR), National Institute of Allergy and Infectious Diseases (NIAID), National Institute of Neurological Disorders and Stroke (NINDS), National Institute of Mental Health (NIMH), National Institute on Drug Abuse (NIDA), National Institute of Nursing Research (NINR), National Cancer Institute (NCI), National Institute on Alcohol Abuse and Alcoholism (NIAAA), National Institute on Deafness and Other Communication Disorders (NIDCD), National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK), and National Institute on Minority Health and Health Disparities (NIMHD) and in coordination and alignment with the research priorities of the National Institutes of Health, Office of AIDS Research (OAR). MWCCS data collection is also supported by UL1-TR000004 (UCSF CTSA), UL1-TR003098 (JHU ICTR), UL1-TR001881 (UCLA CTSI), P30-AI-050409 (Atlanta CFAR), P30-AI-073961 (Miami CFAR), P30-AI-050410 (UNC CFAR), P30-AI-027767 (UAB CFAR), and P30-MH-116867 (Miami CHARM).

The authors have no conflicts of interest to disclose.

A.A.C. conceptualized the study, drafted the manuscript, assisted with the analysis, and revised the paper. J.P.J., L.S., E.A.F., and T.B.N. assisted with the analysis and revised the paper. M.H.C., T.E.W., AR, A.A.A., S.K., A.N.S., M.A.F., A.A., J.M.T., and P.C.T. contributed to study design and data collection and revised the paper. S.D.W. conceptualized the study, contributed to study design and data collection, and revised the paper. All authors approved the final version of the paper.

Correspondence to: Amy A. Conroy, PhD, MPH, Department of Medicine, Center for AIDS Prevention Studies, University of California San Francisco, 550 16th Street, 3rd Floor, San Francisco, CA 94143 (e-mail: amy.conroy@ucsf.edu).

Copyright © 2021 Wolters Kluwer Health, Inc. All rights reserved.

GBV and engagement in care and whether associations were mediated by depression, generalized anxiety disorder (GAD), and posttraumatic stress disorder (PTSD) and moderated by food insecurity.

**Results:** GBV was associated with higher odds of suboptimal adherence (adjusted odds ratio: 1.88; 95% confidence interval: 1.24 to 2.87) and missed appointments (adjusted odds ratio: 1.76; 95% confidence interval: 1.16 to 2.67). The association between GBV and adherence was mediated by depressive symptoms, GAD, and PTSD, accounting for 29.7%, 15.0%, and 16.5%, respectively, of the total association. The association between GBV and missed appointments was mediated by depression and GAD, but not PTSD, with corresponding figures of 25.2% and 19.7%. Associations did not differ by food insecurity.

**Conclusions:** GBV is associated with suboptimal engagement in care, which may be explained by mental health. Interventions should address women's mental health needs, regardless of food insecurity, when improving engagement in HIV care.

**Key Words:** gender-based violence, mental health, food insecurity, antiretroviral adherence

(*J Acquir Immune Defic Syndr* 2022;89:151–158)

## INTRODUCTION

Despite having effective antiretroviral therapy (ART), engagement in HIV care and treatment remains suboptimal among US women living with HIV (WLHIV).<sup>1,2</sup> Gender-based violence (GBV) is a critical barrier to engagement in HIV care among WLHIV. GBV refers to violence perpetrated against an individual or a group on the basis of gender and can take the form of physical, sexual, or emotional abuse by intimate partners, family members, community members, and others.<sup>3,4</sup> The prevalence of intimate partner violence (IPV), a common type of GBV,<sup>5</sup> among WLHIV in the United States is 55%, which is more than double the rate in the general population.<sup>6,7</sup> GBV is associated with poorer HIV clinical outcomes, including detectable viral load, lower CD4 counts, and ART failure.<sup>8–13</sup> A meta-analysis found that IPV had a larger effect on ART adherence than stigma, financial constraints, and pill burden.<sup>14</sup>

Women who experience GBV may miss pills or HIV care appointments for multiple reasons; for example, if they have not disclosed their HIV status to a partner, if an abusive partner prevents them from taking medications,<sup>15–17</sup> or due to feelings of hopelessness.<sup>18,19</sup> GBV is associated with depression and post-traumatic stress disorder (PTSD),<sup>20–23</sup> and poor mental health is linked with nonadherence to ART.<sup>24–27</sup> Thus, one of the most plausible mediating pathways is that GBV leads to poorer mental health, which, in turn, leads to poor engagement in HIV care and treatment and poor clinical outcomes (Fig. 1).<sup>8,14</sup>

It is also plausible that poverty and structural barriers such as food insecurity may worsen the impact of GBV on engagement in care (Fig. 1). Nearly half of WLHIV in the United States are food insecure,<sup>28</sup> defined as having limited access to food and ability to acquire food.<sup>29</sup> Previous research has also shown strong associations between GBV and food insecurity<sup>30–33</sup> and between food insecurity and suboptimal

adherence.<sup>34–36</sup> Women who report food insecurity may miss taking their HIV medications to prevent the side effects of taking ART on an empty stomach and when making trade-offs between spending limited resources on food instead of on transport to pickup medications.<sup>35</sup> Women who are food insecure are more vulnerable to the negative impacts of GBV. In a qualitative study among WLHIV in the United States, the stress of being food insecure, coupled with experiences of violence, led to a low appetite and made it difficult to take HIV medications with food.<sup>37</sup>

To effectively reduce the impact of GBV on engagement in care, several gaps in knowledge must be addressed. Although the mental health pathways linking GBV to suboptimal engagement in care have been described,<sup>8,14,37,38</sup> few studies have explicitly tested whether poor mental health mediates the association between GBV and engagement in care. Moreover, qualitative research has highlighted a complex relationship between food insecurity, GBV, and HIV care engagement,<sup>37</sup> yet no studies have tested whether food insecurity moderates the association between GBV and adherence—which could inform who should be prioritized for interventions.

This study leveraged data from the Women's Inter-agency HIV Study (WIHS) to quantify the effects of GBV (including physical, sexual, and psychological violence) on suboptimal engagement in HIV care and treatment. We also sought to identify the mental health pathways involved, including depressive symptoms, generalized anxiety disorder (GAD), and PTSD. Mental health disorders affect many women who have experienced GBV, which typically manifest as symptoms of depression, anxiety, and PTSD.<sup>39,40</sup> Given our previous research<sup>30,31</sup> and the evidence base suggesting the importance of food insecurity for HIV treatment outcomes for women,<sup>34–36</sup> we assessed whether the association between GBV and engagement in care differs by food insecurity status. We hypothesized that: (1) GBV will be associated with poorer engagement in care (defined as suboptimal adherence to ART and missed clinic appointments) directly and indirectly through mental health and (2) the association between GBV and engagement in care will be greater for women who are food insecure vs. women who are food secure.

## MATERIALS AND METHODS

### Study Population and Procedures

We used longitudinal data from WIHS, a multisite, prospective cohort study of US women at risk of or living with HIV (now continuing as the MACS/WIHS Combined Cohort Study). Study procedures and eligibility criteria have been described elsewhere.<sup>41</sup> In brief, WIHS participants were recruited to represent the demographic profiles of WLHIV in the United States across 10 cities: Bronx, NY; Brooklyn, NY; Washington, DC; Chicago, IL; San Francisco, CA; Chapel Hill, NC; Miami, FL; Birmingham, AL; Jackson, MS; and Atlanta, GA. Women completed interviewer-administered questionnaires every 6 months on demographic characteristics, mental health, violence, psychosocial factors, and engagement in HIV care and treatment and had a brief clinical examination with laboratory tests. From April 2013 to April 2016 (visits

38–43), a module on food insecurity was added to the questionnaire as part of the WIHS Food Insecurity Substudy. Food insecurity data were collected over 6 study visits every 6 months. Participants provided written informed consent and were compensated for participation. The protocol was approved by institutional review boards at all sites in addition to the WIHS Executive Committee.

**Measures**

**Primary Explanatory Variable**


GBV was any sexual, physical, or psychological violence in the past 6 months. Sexual violence was assessed with the question, “Since your (month) study visit, has anyone pressured or forced you to have sexual contact? By sexual contact, I mean them touching your sexual parts, you touching their sexual parts, or sexual intercourse.” Response options were yes, no, do not know, and declined to answer. Physical violence was assessed with the question, “Since your (month) study visit, have you experienced serious physical violence (physical harm by another person)? By that I mean were you ever hurt by a person using an object or were you ever slapped, hit, punched, kicked.” The 2 questions on sexual and physical violence were in reference to “any person,” which could include both partners and non-partners. The questions on psychological violence were in reference to a “current or previous” partner. This was assessed with 7 items (yes/no) such as whether a partner “threatened to hurt you or kill you, “prevented you from leaving or entering the house,” and “prevented you from seeing your friends.” Women who responded yes to any items were coded as having experienced psychological violence.

**Mental Health Mediators**

Depressive symptoms was a continuous variable measured using the Center for Epidemiological Studies Depression scale,<sup>42</sup> collected among all WIHS women during the Food Insecurity Substudy from April 2013 to April 2016 (visits 38–43). Anxiety was a continuous variable measured using the Generalized Anxiety Disorder (GAD)-7 scale,<sup>43</sup> collected from October 2013 to April 2016 (visits 39–43). PTSD symptoms was a continuous variable measured using the PTSD Checklist-Specific (PCL-S),<sup>44</sup> collected as part of the Neurocognition Substudy among a subset of women from April 2013 to April 2015 (visits 38–41). To reduce respondent burden, 25% of women answered the PCL-S items at one of the 4 visits with the aim that 100% of women would have a single measurement for PTSD across the 4-visit period. We created a cross-sectional data set by collapsing the data collected across the 4 visits for analyses examining PTSD as a mediator. GBV, engagement in care, and other variables were captured at the same visit that PTSD was assessed.

**Food Insecurity as a Moderator**

Food insecurity was measured by the US Household Food Security Survey Module (HFSSM),<sup>45</sup> consisting of 18 items asking about the past 6 months (eg, “We worried whether our food would run out before we got money to buy more.”).<sup>46</sup> Using the HFSSM scoring algorithm, we catego-



**FIGURE 1.** Mechanisms linking GBV with engagement in HIV care and treatment.

rized individuals as having high food security, marginal food security (some uncertainty about food supplies, but little to no indications of change in diet or food intake), low food security (reduced quality, variety, or desirability of diet, but little or no indication of reduced food intake), or very low food security (multiple indications of disrupted eating patterns and reduced food intake). Marginal, low, and very low food security represent increasing severity of food insecurity. The HFSSM scale demonstrated high reliability (alpha = 0.91) in WIHS.

**Primary Outcome Variables (Engagement in HIV Care and Treatment)**

Suboptimal adherence to ART was assessed by self-report of how often participants took HIV medications as prescribed over the past 6 months (100% of the time; 95%–99% of the time; 75%–94% of the time; <75% of the time; and I have not taken any of my prescribed medications) and was captured from April 2013 to April 2016. Because a cutoff of 95% or higher represents an optimal level of adherence,<sup>47,48</sup> we dichotomized suboptimal adherence as reporting less than 95% adherence. Missed HIV care appointments were captured from October 2013 to April 2016 (visits 39–43) and was defined as having missed at least one scheduled HIV care appointment in the past 6 months that was not rescheduled (yes/no).

**Covariates**

Based on previous research in WIHS,<sup>28,31</sup> multivariable models adjusted for annual household income (≤\$12,000, \$12,001–\$24,000, \$24,001–\$36,000, \$36,001–\$75,000, and >\$75,000), race/ethnicity (non-Hispanic White, Hispanic, non-Hispanic Black/African American, or other), age (continuous variable), education (<high school degree or equivalent vs. ≥high school degree), and marital status (married/cohabitating vs. unmarried), and time on ART in years.

**Analysis**

We computed sample descriptive characteristics at the first visit of the WIHS Food Insecurity Substudy. We examined the associations between current GBV reported in the past 6 months with suboptimal adherence to ART and missed HIV care appointments in the past 6 months. To

Downloaded from http://journals.lww.com/jaids by BhdMf5ePfkKav1zEum11QIN4+hKjLhEzjgbsH04XMM0hCwWCX1A VhYyQp/1QIH3D00DFRy7T7vSF4C3VC1y0abgQZxdwMfKZB7vys= on 07/12/2023

assess the relative importance of current vs. previous GBV as explanatory variables, we included a lagged variable for GBV in the initial models. Current GBV refers to the same visit in which adherence was assessed, whereas previous GBV refers to a lagged visit 6 months before the current visit. We found that previous GBV was not associated with current adherence to ART or missed appointments, and thus, our final models included only current GBV as the primary explanatory variable. To take full advantage of the longitudinal data, analysis models included data from all visits. We used 2-level logistic regression to model the association between GBV and suboptimal engagement in care or treatment, with women as a random effect, adjusting for covariates. We used random rather than fixed effects because we expected that time-invariant differences between individuals could influence women's engagement in care, and we wanted to directly estimate the association of these variables with our outcomes. Statistical interaction between GBV and food security was assessed using the multiplicative scale by including an interaction term in the multivariable models. The final multivariable analysis used longitudinal data from approximately 1600 women at the baseline visit. All analyses were performed using Stata 15.

Based on our conceptual model and the mediation method used by Weiser et al within WIHS,<sup>28</sup> we estimated the mental health pathways through which GBV may affect engagement in care or treatment. We calculated the direct and indirect effects for each mental health pathway by taking the products of the regression coefficients of each leg of the path from fully adjusted mediation models that included one mental health mediator at a time.<sup>49</sup> Indirect effects were calculated only for paths in which each leg was statistically significant ( $P < 0.05$ ). We confirmed the assumption that each mental health mediator did not have a statistical interaction with GBV for each outcome.<sup>49</sup>

## RESULTS

### Sample Characteristics

At the baseline visit ( $N = 1732$ ), the mean age was 48 years, 67% had a high school education or less, 72% was non-Hispanic Black, and 28% reported being married (Table 1). Across their lifetime, 62% of women reported ever experiencing GBV. At the baseline visit, 5.2% of women experienced sexual or physical violence or psychological violence in the past 6 months. Nearly half of women (44%) experienced any food insecurity (defined as marginal, low, or very low food security). The prevalence of depression, GAD, and PTSD symptoms using established cutoffs for each scale was 35%, 17%, and 14%, respectively. Two-thirds of women were virally suppressed (66%), 17% reported suboptimal adherence, and 14% missed at least one care appointment in the past 6 months.

### Suboptimal Adherence to ART

In the unadjusted models, current experience of GBV was associated with suboptimal adherence to ART in the past

6 months (Table 2). Previous GBV (ie, lagged GBV) was not associated with suboptimal adherence to ART, as noted in the Methods section. Depressive symptoms, GAD, and PTSD symptoms were also associated with suboptimal adherence to ART in the unadjusted models. In the adjusted models, current GBV remained associated with 1.88 higher odds of suboptimal adherence [95% confidence interval (CI): 1.24 to 2.87]. Adjusted models also showed that for each one-unit increase in depression score, there was 1.04 higher odds of suboptimal adherence (95% CI: 1.03 to 1.05). In addition, for each one-unit higher GAD score, there was 1.06 higher odds of suboptimal adherence (95% CI: 1.04 to 1.08), and for each one-unit increase in PTSD score, there was 1.40 higher odds of suboptimal adherence (95% CI: 1.13 to 1.74). Food insecurity did not modify the association between GBV and suboptimal adherence (not included in models in Table 2).

### Missed Care Appointments

In the unadjusted models, current GBV was associated with missing at least one HIV care appointment in the past 6 months (Table 2). Previous GBV was not associated with missed care appointments. Depression and GAD symptoms were associated with missed care appointments in the unadjusted models. In the adjusted models, the association of current GBV with missed care appointment was attenuated and was associated with 1.76 higher odds of a missed care appointment (95% CI: 1.16 to 2.67). Adjusted models also showed that for each one-unit increase in depression score, there was 1.03 higher odds of a missed care appointment (95% CI: 1.01 to 1.03). In addition, for each one-unit increase in GAD score, there was 1.05 higher odds of a missed care appointment (95% CI: 1.03 to 1.07). Food insecurity did not modify the association between GBV and missed care appointments.

### Mediation Analyses

Table 3 lists the total, direct, and indirect effects for each mediation model by the 2 outcomes variables. The adjusted mediation models for suboptimal adherence showed that depression, GAD, and PTSD symptoms accounted for 29.7%, 15.0%, and 16.5% of the total association of GBV with suboptimal adherence, respectively (Table 3). The adjusted mediation models for missed care appointment showed that depression and GAD symptoms accounted for 25.2% and 19.7% of the total association of GBV with missed care appointments, respectively.

## DISCUSSION

In this longitudinal study with more than 1700 WLHIV, GBV was independently associated with suboptimal adherence to ART and at least one missed care visit. These findings are consistent with 2 reviews, which found that women who experience violence are less likely to adhere to ART and achieve viral suppression.<sup>14,50</sup> Most studies have used cross-sectional designs and did not use large, national samples of US WLHIV, other than a WIHS study that found that women

**TABLE 1.** Descriptive Statistics for the Study Sample (N = 1732)

Variable	N (%) or Mean (SD)
Age	47.8 (9.1)
Race/ethnicity	
Non-Hispanic White	178 (10.8%)
Hispanic	244 (14.7%)
Black	1186 (71.6%)
Other	48 (2.9%)
Annual household income	
\$12000 or less	853 (53.7%)
\$12001–24000	354 (22.3%)
\$24001–36000	174 (11.0%)
\$36001–75000	138 (8.7%)
\$75001 or higher	70 (4.4%)
High school education or higher	1111 (67.1%)
Married	489 (28.2%)
Any lifetime GBV	1021 (61.6%)
Any recent GBV	85 (5.2%)
Food insecurity status	
High food security	923 (56.7%)
Marginal food security	252 (15.5%)
Low food security	242 (14.9%)
Very low food security	212 (13.0%)
Depressive symptoms (yes/no)	579 (35.3%)
Generalized anxiety disorder (yes/no)	183 (16.9%)
PTSD (yes/no)	126 (13.7%)
Suboptimal adherence to ART (<95%)	302 (17.4%)
Undetectable viral load	1060 (66.1%)
Missed 1+ clinic appointments	136 (14.2%)
Length of time on ART (yr, median, IQR)	7.8 (3.0–14.5)

who experienced any physical or sexual abuse were less likely to be taking ART.<sup>51</sup> To assess the timing of GBV in relation to engagement in care, we evaluated whether previous and current GBV predicts engagement in care and found that only current GBV was predictive and, therefore,

retained current GBV in our mediation models. Although we cannot rule out reverse causality, one explanation may be that acute experiences of GBV have more immediate effects on missed pills and appointments as opposed to previous experiences of GBV that occurred well before the event of nonadherence.

We also extend previous research by explicitly testing for mental health mediators. Our study suggests that GBV affects engagement in care through depression, GAD, and PTSD symptoms. Depression and GAD were mediators for both adherence to ART and appointments, although PTSD only mediated the association with ART adherence. The role of PTSD should not be discounted, however, because PTSD measures were collected in a smaller subset, and thus, it is possible we would have found a mediating relationship with appointment attendance in a larger sample. Overall, these findings substantiate qualitative studies describing how GBV triggers emotional distress and depressive symptoms, causing forgetfulness to take medications and intentional missing of pills and appointments.<sup>15,17,18,52</sup> Although previous research has not identified the exact time of when symptoms of depression (or anxiety or PTSD) develop after a traumatic event, most studies examine a period of 3–6 months for observing changes in mental health in response to a stressor. A seminal study found that most severe events lead rapidly to depression, occurring in the first 3 weeks after a negative event, and another study found that most of the life events associated with major depression onset occurred within the first month after the event.<sup>53,54</sup> Although certainly GBV can have long-term impacts on mental health, the shorter-term associations that we found by assessing GBV, mental health, and adherence within a 6-month period may indicate a pattern of shorter-term impacts of GBV.

Contrary to our hypothesis, we did not find that food insecurity moderates the association between GBV and engagement in care. Previous work within WIHS has shown that GBV and food insecurity are associated and that food insecurity is associated with worse HIV treatment outcomes.<sup>28,31</sup> Approximately two-thirds of women who

**TABLE 2.** Logistic Regression Models Examining Associations Between GBV and Engagement in HIV Care and Treatment in the Women’s Interagency HIV Study: United States, 2013–2016

Variable	Unique Women (Person Visits)	Odds Ratio (95% CI)	P	Unique Women (Person Visits)	Adjusted Odds Ratio (95% CI)	P
Analysis 1: suboptimal adherence to ART						
Explanatory variable GBV	1689 (7438)	1.86 (1.23 to 2.82)	1665 (7203)	1665 (7203)	1.88 (1.24 to 2.87)	0.003
Mediators						
Depressive symptoms	1690 (7434)	1.04 (1.03 to 1.05)	<0.001	1666 (7197)	1.04 (1.03 to 1.05)	<0.001
GAD symptoms	1637 (5987)	1.05 (1.03 to 1.08)	<0.001	1612 (5806)	1.06 (1.04 to 1.08)	<0.001
PTSD symptoms	840	1.31 (1.08 to 1.59)	0.006	803	1.40 (1.13 to 1.74)	0.002
Analysis 2: missed HIV care appointments						
Explanatory variable GBV	1732 (6430)	1.97 (1.30 to 3.00)	<0.001	1706 (6231)	1.76 (1.16 to 2.67)	0.008
Mediators						
Depressive symptoms	1732 (6424)	1.03 (1.02 to 1.04)	<0.001	1706 (6224)	1.03 (1.01 to 1.03)	<0.001
GAD symptoms	1735 (6466)	1.06 (1.04 to 1.07)	<0.001	1709 (6256)	1.05 (1.03 to 1.07)	<0.001
PTSD symptoms	899	1.19 (0.94 to 1.50)	0.140	8wp66	1.03 (0.80 to 1.33)	0.830

Adjusted models controlled for age at visit, employment status, income, race or ethnicity, education, marital status, and time on ART.

Downloaded from http://journals.lww.com/jaids by BHD/M/5e/P/H/Kav12Eoun11Q/N4+hK/LhEZgbsH04XMM0hC/wC/1A VhYQp/llQH/D3D0DQFR/y7TvsF4C3/VC1y0abgqZXdwnfKZBYtws= on 07/12/2023

**TABLE 3.** Mediating Effects of Mental Health in the Association Between GBV and Engagement in HIV Care and Treatment

Mediator Models	Total Association	Direct Association	Indirect Association	Proportion Mediated
Analysis 1: association between GBV and suboptimal adherence to ART				
Depressive symptoms	0.59	0.41	0.18	29.73%
Generalized anxiety disorder	0.74	0.63	0.11	15.03%
PTSD	1.08	0.90	0.18	16.54%
Analysis 2: association between GBV and missed HIV care appointments				
Depressive symptoms	0.47	0.35	0.12	25.23%
Generalized anxiety disorder	0.54	0.43	0.11	19.68%

Separate mediator models were run for each mediating variable. All models controlled for age at visit, employment status, income, race or ethnicity, education, marital status, and time on ART.

reported recent GBV had food insecurity; however, the impact of GBV on care and treatment was not exacerbated by food insecurity. It is possible that although food insecurity is independently linked to both violence and engagement in care, it plays less of a role in the association between the 2. An additional explanation, which we did not test, could be that food insecurity leads to violence, which then leads to nonadherence through serial mediation. Similar relationships should be examined for other dimensions of material need insecurities (eg, housing instability, poor access to health care, etc.) that may play an important role.<sup>55</sup>

Our findings have important implications for programming. First, our findings suggest that it may be insufficient to reduce women's exposure to GBV when mental health is one of the main mechanisms through which ART adherence is affected. For example, it may be not be enough to refer women living with HIV to domestic violence shelters without simultaneously addressing their mental health. Existing interventions, most of which have focused on trauma around childhood sexual abuse, could be adapted to address GBV,<sup>56–58</sup> mental health, and engagement in care for WLHIV. In addition, victims of violence could be screened for mental health symptoms and referred to psychological services such as trauma-focused cognitive behavioral therapy, which could be integrated into HIV clinical care.<sup>59</sup> Based on our findings suggesting that exposure to violence may have short-term effects on the development of mental health symptoms, it would be prudent to intervene quickly on mental health before conditions worsen (eg, immediate referrals and linkage into mental health services). Of course, programs that focus on the primary prevention of violence are also needed to prevent the onset of mental health symptoms and subsequent nonadherence. Finally, based on our finding that food insecurity did not worsen the association between GBV and engagement in care, which could have signaled the need to prioritize who are food insecure, we recommend that

violence interventions target all WLHIV. However, interventions still need to consider the importance of food insecurity for suboptimal adherence among WLHIV.<sup>60</sup>

Several limitations are noteworthy. Although the use of longitudinal data with lagged variables and mediators strengthens the plausibility of a causal link, it is still possible that poor adherence or mental health leads to GBV. Violence and mental health have a bidirectional relationship as shown through longitudinal studies,<sup>61</sup> although this evidence is scarce. GBV and mental health could operate in a vicious cycle, mutually affecting the other, to affect care engagement. Although evidence consistently shows that violence leads to poor mental health, it is possible that poor mental health could increase the risk of violence or conflict in interpersonal relationships. A second related limitation is our inability to show temporal ordering of events. We included all women across all visits in our analysis rather than selecting 3 time points to capture the temporal ordering of GBV, mental health, and adherence. This would have significantly reduced the number of observations in the analysis and the time gap between violence and adherence would be too long (eg, 1 year) for violence to affect mental health and then adherence, which we believed would occur in a shorter period. Indeed, we found that current, but not previous, GBV is associated with poor care engagement. Future studies could capture smaller intervals such as every 3 months or daily to parse out the temporal ordering of events. Third, our measure of GBV included both IPV and violence from nonpartners. Given that IPV is the most common form, it is likely that our measure of GBV is mostly capturing IPV; however, future studies should separate out the 2 types of violence to assess the same patterns hold. Finally, we relied on self-reported measures of violence and adherence, subject to recall and social desirability bias.

## CONCLUSIONS

In summary, the results provide strong evidence that GBV is associated with suboptimal engagement in HIV care and treatment. This may occur through mental health pathways, including depression, GAD, and PTSD symptoms. Food insecurity did not modify the associations of GBV on adherence and engagement in care. Interventions for WLHIV should strive to reduce experiences of GBV while also addressing women's mental health to sustain progress toward improving HIV clinical outcomes and prevent the spread of HIV.

## ACKNOWLEDGMENTS

*The authors thank the participants in the WIHS for their contributions to the study overall and specifically to this research.*

## REFERENCES

1. Crepez N, Tang T, Marks G, et al. Viral suppression patterns among persons in the United States with diagnosed HIV infection in 2014. *Ann Intern Med.* 2017;167:446–447.
2. Crepez N, Tang T, Marks G, et al. Durable viral suppression and transmission risk potential among persons with diagnosed HIV infection: United States, 2012–2013. *Clin Infect Dis.* 2016;63:976–983.

3. USAID. *United States Strategy to Prevent and Respond to Gender-Based Violence Globally: 2016 Update*. Washington, DC: USAID, 2016.
4. United Nations. *Declaration on the Elimination of Violence against Women*. New York, NY: United Nations, 1993.
5. Watts C, Zimmerman C. Violence against women: global scope and magnitude. *Lancet*. 2002;359:1232–1237.
6. Machtinger E, Wilson T, Haberer JE, et al. Psychological trauma and PTSD in HIV-positive women: a meta-analysis. *AIDS Behav*. 2012;16:2091–2100.
7. Smith SG, Basile KC, Gilbert LK, et al. *National Intimate Partner and Sexual Violence Survey (NISVS): 2010–2012 State Report*. Atlanta, GE: National Center for Injury Prevention and Control; 2017.
8. Schafer KR, Brant J, Gupta S, et al. Intimate partner violence: a predictor of worse HIV outcomes and engagement in care. *AIDS Patient Care STDS*. 2012;26:356–365.
9. Machtinger E, Haberer J, Wilson T, et al. Recent trauma is associated with antiretroviral failure and HIV transmission risk behavior among HIV-positive women and female-identified transgenders. *AIDS Behav*. 2012;16:2160–2170.
10. Espino SR, Fletcher J, Gonzalez M, et al. Violence screening and viral load suppression among HIV-positive women of color. *AIDS Patient Care STDS*. 2015;29:S36–S41.
11. Rose RC, House AS, Stepleman LM. Intimate partner violence and its effects on the health of African American HIV-positive women. *Psychol Trauma Theor Res Pract Pol*. 2010;2:311.
12. Trimble DD, Nava A, McFarlane J. Intimate partner violence and antiretroviral adherence among women receiving care in an urban Southeastern Texas HIV clinic. *J Assoc Nurses AIDS Care*. 2013;24:331–340.
13. Siemieniuk RA, Krentz HB, Miller P, et al. The clinical implications of high rates of intimate partner violence against HIV-positive women. *J Acquir Immune Defic Syndr*. 2013;64:32–38.
14. Hatcher AM, Smout EM, Turan JM, et al. Intimate partner violence and engagement in HIV care and treatment among women: a systematic review and meta-analysis. *AIDS*. 2015;29:2183–2194.
15. Lichtenstein B. Domestic violence in barriers to health care for HIV-positive women. *AIDS Patient Care STDS*. 2006;20:122–132.
16. Maeri I, El Ayadi A, Getahun M, et al. “How can I tell?” Consequences of HIV status disclosure among couples in eastern African communities in the context of an ongoing HIV “test-and-treat” trial. *AIDS Care*. 2016;28:59–66.
17. Orza L, Bass E, Bell E, et al. In women’s eyes: key barriers to Women’s access to HIV treatment and a rights-based approach to their sustained well-being. *Health Hum Rights*. 2017;19:155.
18. Conroy A, Leddy A, Johnson M, et al. “I told her this is your life”: relationship dynamics, partner support and adherence to antiretroviral therapy among South African couples. *Cult Health Sex*. 2017;19:1239–1253.
19. Hatcher A, Stöckl H, Christofides N, et al. Mechanisms linking intimate partner violence and prevention of mother-to-child transmission of HIV: a qualitative study in South Africa. *Soc Sci Med*. 2016;168:130–139.
20. Conroy A, Tuthill E, Neilands T, et al. *A Dyadic Investigation of Relationship Dynamics and Depression in HIV-Infected Couples from Malawi*. Poster presented at the International AIDS Conference 23–27 July 2018. Amsterdam, Netherlands; 2018.
21. Pico-Alfonso MA, Garcia-Linares MI, Celda-Navarro N, et al. The impact of physical, psychological, and sexual intimate male partner violence on women’s mental health: depressive symptoms, posttraumatic stress disorder, state anxiety, and suicide. *J Womens Health*. 2006;15:599–611.
22. Martinez J, Hosek SG, Carleton RA. Screening and assessing violence and mental health disorders in a cohort of inner city HIV-positive youth between 1998–2006. *AIDS Patient Care STDS*. 2009;23:469–475.
23. Ellsberg M, Jansen HA, Heise L, et al. Intimate partner violence and women’s physical and mental health in the WHO multi-country study on women’s health and domestic violence: an observational study. *Lancet*. 2008;371:1165–1172.
24. Hahn JA, Samet JH. Alcohol and HIV disease progression: weighing the evidence. *Curr HIV/AIDS Rep*. 2010;7:226–233.
25. Hinkin CH, Hardy DJ, Mason KI, et al. Medication adherence in HIV-infected adults: effect of patient age, cognitive status, and substance abuse. *AIDS*. 2004;18(suppl 1):S19.
26. Arnsten JH, Demas PA, Grant RW, et al. Impact of active drug use on antiretroviral therapy adherence and viral suppression in HIV-infected drug users. *J Gen Intern Med*. 2002;17:377–381.
27. Lucas GM, Gebo KA, Chaisson RE, et al. Longitudinal assessment of the effects of drug and alcohol abuse on HIV-1 treatment outcomes in an urban clinic. *AIDS*. 2002;16:767–774.
28. Weiser SD, Sheira LA, Palar K, et al. Mechanisms from food insecurity to worse HIV treatment outcomes in US women living with HIV. *AIDS Patient Care STDS*. 2020;34:425–435.
29. National Research Council. *Food Insecurity and Hunger in the United States: An Assessment of the Measure*. Washington, DC: National Academies Press; 2006.
30. Diamond-Smith N, Conroy AA, Tsai AC, et al. Food insecurity and intimate partner violence among married women in Nepal. *J Glob Health*. 2019;9:010412.
31. Conroy AA, Cohen M, Frongillo E, et al. Food insecurity and violence in a prospective cohort of U.S. women at risk or living with HIV. *PLoS One*. 2019;14:e0213365.
32. Hatcher AM, Stöckl H, McBride RS, et al. Pathways from food insecurity to intimate partner violence perpetration among peri-urban men in South Africa. *Am J Prev Med*. 2019;56:765–772.
33. Hatcher AM, Weiser SD, Cohen CR, et al. Food insecurity and intimate partner violence among HIV-positive individuals in rural Kenya. *Am J Prev Med*. 2021;60:563–568.
34. Weiser SD, Palar K, Frongillo EA, et al. Longitudinal assessment of associations between food insecurity, antiretroviral adherence and HIV treatment outcomes in rural Uganda. *AIDS*. 2014;28:115–120.
35. Young S, Wheeler AC, McCoy SI, et al. A review of the role of food insecurity in adherence to care and treatment among adult and pediatric populations living with HIV and AIDS. *AIDS Behav*. 2014;18:505–515.
36. Palar K, Sheira LA, Frongillo EA, et al. Longitudinal relationship between food insecurity, engagement in care, and ART adherence among US women living with HIV. *AIDS*. 2013;28:35.
37. Leddy AM, Zakaras JM, Shieh J, et al. Intersections of food insecurity, violence, poor mental health and substance use among US women living with and at risk for HIV: evidence of a syndemic in need of attention. *PLoS One*. 2021;16:e0252338.
38. LeGrand S, Reif S, Sullivan K, et al. A review of recent literature on trauma among individuals living with HIV. *Curr HIV/AIDS Rep*. 2015;12:397–405.
39. Rees S, Silove D, Chey T, et al. Lifetime prevalence of gender-based violence in women and the relationship with mental disorders and psychosocial function. *JAMA*. 2011;306:513–521.
40. Walsh K, Keyes KM, Koenen KC, et al. Lifetime prevalence of gender-based violence in US women: associations with mood/anxiety and substance use disorders. *J Psychiatr Res*. 2015;62:7–13.
41. Adimora AA, Ramirez C, Benning L, et al. Cohort profile: the women’s interagency HIV study (WIHS). *Int J Epidemiol*. 2018;47:393–394i.
42. Radloff LS. The CES-D scale a self-report depression scale for research in the general population. *Appl Psychol Meas*. 1977;1:385–401.
43. Spitzer RL, Kroenke K, Williams JB, et al. A brief measure for assessing generalized anxiety disorder: the GAD-7. *Arch Intern Med*. 2006;166:1092–1097.
44. Ruggiero KJ, Del Ben K, Scotti JR, et al. Psychometric properties of the PTSD checklist—civilian version. *J Trauma Stress*. 2003;16:495–502.
45. Bickel G, Nord M, Price C, et al. *Guide to Measuring Household Food Security*. Alexandria, VA: Food and Nutrition Service, USDA; 2000.
46. Frongillo EA. Validation of measures of food insecurity and hunger. *J Nutr*. 1999;129:506S–509S.
47. Kelso GA, Cohen MH, Weber KM, et al. Critical consciousness, racial and gender discrimination, and HIV disease markers in African American women with HIV. *AIDS Behav*. 2014;18:1237–1246.
48. Dale S, Cohen M, Weber K, et al. Abuse and resilience in relation to HAART medication adherence and HIV viral load among women with HIV in the United States. *AIDS Patient Care STDS*. 2014;28:136–143.
49. VanderWeele T. *Explanation in Causal Inference: Methods for Mediation and Interaction*. Oxford, UK: Oxford University Press; 2015.
50. Leddy AM, Weiss E, Yam E, et al. Gender-based violence and engagement in biomedical HIV prevention, care and treatment: a scoping review. *BMC Public Health*. 2019;19:897.

51. Cohen MH, Cook JA, Grey D, et al. Medically eligible women who do not use HAART: the importance of abuse, drug use, and race. *Am J Public Health*. 2004;94:1147–1151.
52. Zunner B, Dworkin SL, Neylan TC, et al. HIV, violence and women: unmet mental health care needs. *J Affect Disord*. 2015;174:619–626.
53. Kendler KS, Karkowski LM, Prescott CA. Stressful life events and major depression: risk period, long-term contextual threat, and diagnostic specificity. *J Nervous Ment Dis*. 1998;186:661–669.
54. Brown G, Harris T. *Social Origins of Depression*. London, UK: Free Press; 1978.
55. Whittle HJ, Leddy AM, Shieh J, et al. Precarity and health: theorizing the intersection of multiple material-need insecurities, stigma, and illness among women in the United States. *Soc Sci Med*. 2020;245:112683.
56. Chin D, Myers HF, Zhang M, et al. Who improved in a trauma intervention for HIV-positive women with child sexual abuse histories? *Psychol Trauma Theor Res Pract Pol*. 2014;6:152.
57. Meade CS, Drabkin AS, Hansen NB, et al. Reductions in alcohol and cocaine use following a group coping intervention for HIV-positive adults with childhood sexual abuse histories. *Addiction*. 2010;105:1942–1951.
58. Sikkema KJ, Hansen NB, Kochman A, et al. Outcomes from a group intervention for coping with HIV/AIDS and childhood sexual abuse: reductions in traumatic stress. *AIDS Behav*. 2007;11:49–60.
59. Marshall KJ, Fowler DN, Walters ML, et al. Interventions that address intimate partner violence and HIV among women: a systematic review. *AIDS Behav*. 2018;22:3244–3263.
60. Leddy AM, Sheira LA, Tamraz B, et al. Food insecurity is associated with lower levels of antiretroviral drug concentrations in hair among a cohort of women living with human immunodeficiency virus in the United States. *Clin Infect Dis*. 2020;71:1517–1523.
61. Oram S, Khalifeh H, Howard LM. Violence against women and mental health. *Lancet Psychiatry*. 2017;4:159–170.