

Mutations in the *CYLD* gene in Brooke–Spiegler Syndrome, Familial Cyndromatosis, and Multiple Familial Trichoepithelioma: Lack of Genotype–Phenotype Correlation

Sarah Bowen,* Melissa Gill,† David A. Lee,* Galen Fisher,‡ Roy G. Geronemus,‡ MariaLuisa Espinel Vazquez,§ and Julide Tok Celebi*

Departments of *Dermatology and †Pathology, Columbia University, New York, New York, USA; ‡The Laser and Skin Surgery Center of New York, New York, New York, USA; §Hospital Comarcal de la Linea, Cadiz, Spain

Brooke–Spiegler syndrome (BSS), familial cyndromatosis (FC), and multiple familial trichoepithelioma (MFT), originally described as distinct entities, share overlapping clinical findings. Patients with BSS are predisposed to multiple skin appendage tumors such as cyndroma, trichoepithelioma, and spiradenoma. FC, however, is characterized by cyndromas and MFT by trichoepitheliomas as the only tumor type. These disorders have recently been associated with mutations in the *CYLD* gene. In this report, we describe three families with BSS, one with FC, and two with MFT phenotypes associated with novel and recurrent mutations in *CYLD*. We provide evidence that these disorders represent phenotypic variation of a single entity and lack genotype–phenotype correlation.

Key words: *CYLD*/genodermatosis/mutation/tumor suppressor
J Invest Dermatol 124:919–920, 2005

Brooke–Spiegler syndrome (BSS, OMIM# 605041) (Brooke, 1892; Spiegler, 1899) is an autosomal dominantly inherited disease characterized by multiple skin appendage tumors. The predominating tumor can be a cyndroma, trichoepithelioma, and/or spiradenoma. Although cyndromas and spiradenomas are sweat gland tumors, trichoepitheliomas show hair follicle differentiation. Therefore, it has been postulated that BSS results from defects in the regulation of putative stem cells of the folliculo–sebaceous–apocrine unit, giving rise to different skin appendage tumors (Fenske *et al*, 2000). Two autosomal dominantly inherited genodermatoses, familial cyndromatosis (FC, OMIM# 132700), and multiple familial trichoepithelioma (MFT, OMIM# 601606), have been originally described as distinct entities. FC, originally known as Ansell–Spiegler cyndromas, is characterized by cyndromas, whereas MFT, originally referred to as Brooke–Fordyce trichoepitheliomas, is characterized by trichoepitheliomas (Welch *et al*, 1968).

The gene for FC was mapped to chromosome 16q12–q13 by linkage analysis (Biggs *et al*, 1995). The *CYLD* gene was discovered by positional cloning and germline mutations were identified in families with this disease (Bignell *et al*, 2000). Loss of heterozygosity at the *CYLD* locus has been found in both inherited and sporadic tumors, implicating *CYLD* as a tumor suppressor gene (Bignell *et al*, 2000). Subsequently, mutations in *CYLD* in patients with BSS phenotype were found (Hu *et al*, 2003). More recently, *CYLD* has been identified as the susceptibility gene for several families with MFT (Salhi *et al*, 2004; Zhang *et al*, 2004; Zheng *et al*, 2004). These findings, taken together

with the observations that features of BSS, FC, and MFT can occur in the same patient or in different patients within a single family (Gerretsen *et al*, 1995), and that a single *CYLD* mutation can be associated with both cyndromas and trichoepitheliomas (Poblete Gutierrez *et al*, 2002), suggest that these syndromes not only share a common genetic basis but represent phenotypic variation of the same disease.

To date, there have been 29 germline mutations described in the *CYLD* gene in families with BSS, FC, and MFT (Bignell *et al*, 2000; Poblete Gutierrez *et al*, 2002; Hu *et al*, 2003; Scheinfeld *et al*, 2003; Salhi *et al*, 2004; Zhang *et al*, 2004; Zheng *et al*, 2004). In this report, we describe three novel and two recurrent mutations in *CYLD* identified in five unrelated families with BSS, FC, and MFT phenotypes.

Results

We evaluated three families with BSS, one with FC, and two with MFT phenotypes (Table S1). The pedigree of each family was consistent with an autosomal dominant mode of inheritance (data not shown). In all families except one, mutations in the *CYLD* gene were identified (Fig S1). These mutations are summarized in Table S1. All of the identified mutations were either nonsense or frameshift mutations resulting in premature stop codons. The case in which a mutation was not found (Cyld-3) showed an MFT phenotype with multiple trichoepitheliomas as the only skin tumor.

Discussion

All of the reported mutations in *CYLD* have been heterozygous mutations with the majority (~90%) predicted

Abbreviations: BSS, Brooke–Spiegler syndrome; FC, familial cyndromatosis; MFT, multiple familial trichoepithelioma

to result in truncated proteins or lead to nonsense mediated mRNA decay. To date, no mutations in the N-terminal region of *CYLD* encoding the first two cytoskeletal-associated-protein-glycine-conserved motifs have been reported (exons 4–8). The absence of mutations in the N-terminal region encoding two CAP-GLY domains emphasizes the importance of these motifs for its function. The mutations, however, are scattered throughout the mid-portion and C-terminal region. Although clustering of mutations in the C-terminal region (exons 16–20) is noted (~60%), this is not limited to one particular domain. As a result, for mutation screening strategies, we recommend starting with exons 16–20 and then pursuing the other parts of the gene.

Of interest, in one of the cases described in this report with the MFT phenotype (Cyld-3), we did not find sequence alterations in the coding sequence and flanking introns of *CYLD*. Although we cannot rule out the possibility of a large germline deletion involving *CYLD* in this case, it is possible that another locus for MFT exists. Evaluation of additional families with MFT will help clarify the presence of genetic heterogeneity.

In addition to tumors of the skin appendages, patients with BSS are also at risk for developing tumors of the major and minor salivary glands. Basal cell adenomas and adenocarcinomas of the parotid glands and minor salivary glands have been reported in association with this disease (Jungehulsing *et al*, 1999). One case described in this report (Cyld-6) demonstrated bilateral parotid gland basal cell adenocarcinomas, exemplifying this association.

We also evaluated the mutations in *CYLD* for genotype-phenotype correlation. To date, there have been six mutations described in association with the MFT phenotype. These mutations located in exons 10, 15–18, and 20 show no particular clustering in one region. Based on the current mutational data in individuals with the BSS, FC, and MFT phenotypes, in addition to the observation that within a single family some individuals may have trichoepitheliomas whereas others have cylindromas, it appears a correlation between genotype and phenotype is lacking.

Materials and Methods

Six unrelated families with BSS, FC, or MFT phenotypes were recruited to this study. The Institutional Review Board at Columbia University approved all the described studies and the study was conducted according to Declaration of Helsinki Guidelines. Blood samples from affected individuals were collected after obtaining informed consent. Genomic DNA was extracted from whole blood using PureGene DNA isolation kit (Gentra Systems, Minneapolis, Minnesota). The 17 coding exons of the *CYLD* gene were amplified by PCR using specific primers. For mutation detection, the PCR products were sequenced using an automated sequencing system (310, Applied Biosystems, Foster City, California).

We appreciate the participation of the patients in the study. We are grateful to Steven Wollinsky, MD, Michael T. Geraghty, MD, and Michele Horner, M.S. for referring their patients. This work was supported by the Dermatology Foundation (J. T. C.).

Supplementary Material

The following material is available from <http://www.blackwellpublishing.com/products/journals/suppmat/JID/JID23688/JID23688.htm>

Figure S1. Clinical findings and mutation analysis.

Table S1. Clinical data and mutation status of the families described in this report

DOI: 10.1111/j.0022-202X.2005.23688.x

Manuscript received October 26, 2004; revised December 7, 2004; accepted for publication January 3, 2005

Address correspondence to: Julide Tok Celebi, MD, Department of Dermatology, Columbia University, 630 West 168th Street, VC-15-202 New York, NY 10032, USA. Email: jt165@columbia.edu

References

- Biggs PJ, Wooster R, Ford D, *et al*: Familial cylindromatosis (turban tumour syndrome) gene localised to chromosome 16q12–q13: Evidence for its role as a tumour suppressor gene. *Nat Genet* 11:441–443, 1995
- Bignell GR, Warren W, Seal S, *et al*: Identification of the familial cylindromatosis tumour-suppressor gene. *Nat Genet* 25:160–165, 2000
- Brooke H: Epithelioma adenoides cysticum. *Br J Dermatol* 4:269, 1892
- Fenske C, Banerjee P, Holden C, Carter N: Brooke–Spiegler syndrome locus assigned to 16q12–q13. *J Invest Dermatol* 114:1057–1058, 2000
- Gerretsen AL, Beemer FA, Deenstra W, Hennekam FA, van Vloten WA: Familial cutaneous cylindromas: Investigations in five generations of a family. *J Am Acad Dermatol* 33:199–206, 1995
- Hu G, Onder M, Gill M, Aksakal B, Oztas M, Gurer MA, Celebi JT: A novel missense mutation in *CYLD* in a family with Brooke–Spiegler syndrome. *J Invest Dermatol* 121:732–734, 2003
- Jungehulsing M, Wagner M, Damm M: Turban tumour with involvement of the parotid gland. *J Laryngol Otol* 113:779–783, 1999
- Poblete Gutierrez P, Eggermann T, Holler D, *et al*: Phenotype diversity in familial cylindromatosis: A frameshift mutation in the tumor suppressor gene *CYLD* underlies different tumors of skin appendages. *J Invest Dermatol* 119:527–531, 2002
- Salhi A, Bornholdt D, Oeffner F, Malik S, Heid E, Happle R, Grzeschik KH: Multiple familial trichoepithelioma caused by mutations in the cylindromatosis tumor suppressor gene. *Cancer Res* 64:5113–5117, 2004
- Scheinfeld N, Hu G, Gill M, Austin C, Celebi JT: Identification of a recurrent mutation in the *CYLD* gene in Brooke–Spiegler syndrome. *Clin Exp Dermatol* 28:539–541, 2003
- Spiegler H: Uber endothelioma der haut. *Arch Derm Syph* 50: 163–176, 1899
- Welch JP, Wells RS, Kerr CB: Ansell–Spiegler cylindromas (turban tumours) and Brooke–Fordyce trichoepitheliomas: Evidence for a single genetic entity. *J Med Genet* 5:29–35, 1968
- Zhang XJ, Liang YH, He PP, *et al*: Identification of the cylindromatosis tumor-suppressor gene responsible for multiple familial trichoepithelioma. *J Invest Dermatol* 122:658–664, 2004
- Zheng G, Hu L, Huang W, *et al*: *CYLD* mutation causes multiple familial trichoepithelioma in three Chinese families. *Hum Mutat* 23:400, 2004