

Table 1. Percent frequency of invasive species collected in eight paired shallow and deep water sites in each of four western New York Finger Lakes: Conesus, Honeoye, Canadice and Hemlock

Common Name	Scientific Name	Lake Conesus		Honeoye		Canadice		Hemlock	
		Shallow	Deep	Shallow	Deep	Shallow	Deep	Shallow	Deep
Zebra Mussel	<i>Dreissena polymorpha</i>	38	50	25	63	50	38	75	75
Quagga Mussel	<i>Dreissena bugensis</i>	25	50	25	50	38	25	25	50
Banded Mystery Snail	<i>Viviparum georgianus</i>	25	13	13	38	0	0	0	0
Chinese Mystery Snail	<i>Cipangopaludna chinensis malleata</i>	0	13	13	13	0	0	0	0
Faucet Snail	<i>Bithynia tentaculata</i>	13	0	0	25	0	0	0	0
Euryhaline Amphipod	<i>Echinogammarus ischnus</i>	0	0	0	0	0	0	0	0
Bloody Red Shrimp	<i>Hemimysis anomala</i>	0	0	0	0	0	0	0	0
Asian Clam	<i>Corbicula fluminea</i>	0	0	0	0	0	0	0	0

V. georgianus				D. polymorpha		
Shallow	Deep			Shallow	Deep	
	25	38		38	50	
	13	13		25	63	
	19.0	25.5		50	38	
	8.5	17.7		75	75	
			Mean	47.0	56.5	
C. c. malleata			STDev	21.3	16.0	
	0	13				
	13	13		D. bugensis		
	6.5	13.0		Shallow	Deep	
	9.2	0.0		25	50	
				25	50	

B. tentaculata

13

0

38 25

25 50

0

25

Mean

28.3 43.8

6.5

12.5

STDev

6.5 12.5

9.2

17.7